



CENTER FOR ROBOTIC & METABOLIC SURGERY

PATIENT PREOPERATIVE INSTRUCTIONS

Medication Instructions Before Surgery

Follow the medication plan given by your surgical and anesthesia team. GLP-1 hold instructions below apply only when your team directs you to stop treatment.

GLP-1 MEDICATIONS | HOLD 1 WEEK (7 DAYS)

When directed by your surgical / anesthesia team; confirm your individual hold period.

Brand name	Medication
Ozempic	Semaglutide
Rybelsus	Semaglutide
Wegovy	Semaglutide
Trulicity	Dulaglutide
Victoza	Liraglutide
Saxenda	Liraglutide
Byetta	Exenatide
Bydureon BCise	Exenatide
Mounjaro*	Tirzepatide
Zepbound*	Tirzepatide
Adlyxin	Lixisenatide

*Tirzepatide is a dual GIP/GLP-1 medication. Also report compounded products.

SGLT2 MEDICATIONS | HOLD AT LEAST 72 HOURS

Steglatro requires a longer hold: at least 96 hours (4 days) before surgery.

Brand name	Medication	Stop before surgery
Invokana	Canagliflozin	72 hours (3 days)
Farxiga	Dapagliflozin	72 hours (3 days)
Jardiance	Empagliflozin	72 hours (3 days)
Steglatro	Ertugliflozin	96 hours (4 days)
Brenzavvy	Bexagliflozin	72 hours (3 days)

Medication / dose: _____

Last dose date: _____ Hold starting: _____

Call your care team if you took a dose during the hold period or have nausea, vomiting, bloating or abdominal pain. If you have diabetes, ask your prescriber about blood sugar management. Follow your separate fasting instructions. Restart medications only as directed by your care team.

Bring your medication list on the day of surgery.